

Integrated Pest Management Program

Department of Plant Science and Landscape Architecture Department of Extension

Update for Greenhouse Growers, Tales from the Field, August 9, 2013
Encouraging Good Roots, Rusts, and Miscanthus Blight
Leanne Pundt, Extension Educator, Greenhouse IPM

Encourage Good Root Development on Poinsettias Now – It's important not to stress young roots by high salt or ec levels, compacted media (due to overwatering) or by not controlling low levels of fungus gnat larvae **early**. Waiting until you see extensive damage (see below), makes it almost impossible for the root system to recover before short days begin. Keeping plants actively growing in August and September by developing a good root system is crucial to having high quality poinsettias.



Figure 1: Fungus gnat feeding damage, Photo by L. Pundt Figure 2: Healthy root system, Photo by R. McGaughey

Many growers report that an application of a biological fungicide such as Root Shield combined with the beneficial nematodes for fungus gnat larvae (Nemasys, Nemashield, Scanmask, Entonem) helps get root systems off to a good start. Continue drench application of beneficial nematodes thru the growing season to prevent fungus gnat feeding so you have a healthy root system.

Beneficial Nematodes: An Easy Way to Begin Using Biological Control in the Greenhouse http://s.uconn.edu/1rs

Rusts – the cool night temperatures and heavy dews are favorable for many foliar diseases including rusts and powdery mildew. For information on chrysanthemum white rust, see http://extension.umass.edu/floriculture/fact-sheets/chrysanthemum-white-rust



Rusts on Ornamental Grasses

Rust diseases are often seen on ornamental grasses especially susceptible varieties of Calamagrostis, (Reed grass), and Panicum (Switchgrass) and occasionally Miscanthus (Maiden grass). However, more frequently, I see Miscanthus Blight on Maiden grass.



Figure 3: Elizabeth Bush, Virginia Polytechnic Institute Figure 4: Miscanthus Blight, Photo by L. Pundt and State University, Bugwood.org

Miscanthus can be affected by a number of different leaf spotting fungi. One of the more noticeable is Miscanthus Blight, caused by the fungus Stagonospora. You will see purplish or rust colored spots and streaks on the leaves, especially the white sections of the variegated leaves. From a distance, with its rusty appearance, it is often mistaken for a rust disease.

Rust Diseases on Ornamental Crops http://s.uconn.edu/1ru

