

Cicada Killer, Giant Ground Hornet, *Sphecius hogardii* (Latreille) and *Sphecius speciosus* (Drury) (Insecta: Hymenoptera: Sphecidae)¹

Lionel A. Stange²

Introduction

Cicada killers, or giant ground hornets, are among the largest wasps in Florida (up to 40 mm in length). They are conspicuous insects, since the males are territorial and will butt or grapple with intruders including other males. There are 22 species of *Sphecius* Dahlbom in the world and all hunt cicadas as far as scientists know. Two of four Nearctic species of *Sphecius* occur in Florida. The females of the common Florida species, *Sphecius speciosus* (Drury), hunt *Tibicen* spp. cicadas and can dig four-foot burrows in the ground with several branches and cells. The provisioning with cicadas is nearly specific to *Sphecius* in the family Sphecidae, but is known in a few other sphecids, such as *Liogorytes joergenseni* (Brethes) from Argentina (Bohart and Stange 1976).

Biology

This biology is based on *Sphecius speciosus*. Lin (1966) observed copulatory habits in these wasps. Usually, copulation was initiated while the wasps were on the ground, with the male grasping the female from behind. Then the wasps would fly out with the female dragging the usually smaller male. Their heads were diametrically opposed while in flight. However, both wasps used their wings in flight, which can last up to 45 minutes.

The fertilized female begins the reproductive cycle by making burrows 0.3 to 1.2 m in length, with numerous branches and cells. The front legs rake the soil, which the female pushes out with her hind legs, resulting in a large dirt mound. The burrows are left open during provisioning and may never be closed. However, the cells are carefully closed when completed. Between one and four cicadas are deposited per cell depending on the size of the adult cicada.

The female paralyzes the host by stinging the membrane at the base of the foreleg and then she carries the cicada venter to venter by dragging or flying to the nest. One egg is placed on the last cicada in each nest. Apparently, the mother wasp can determine the sex of the egg and places the female egg with the more abundant provisions. Lin and Michener (1972) recorded up to four females of *Sphecius speciosus* provisioning a single nest simultaneously, which may indicate a step toward sociality.

Key to Species of *Sphecius* in Florida

1. Body predominately rust-orange color with mostly black antennae and last few segments of the abdomen black
Sphecius hogardii (Latreille)

1. This document is EENY295, one of a series of the Department of Entomology and Nematology, UF/IFAS Extension. Original publication date June 2003. Revised August 2012 and November 2015. Reviewed October 2018. Visit the EDIS website at <http://edis.ifas.ufl.edu>. This document is also available on the Featured Creatures website at <http://entnemdept.ifas.ufl.edu/creatures/>.

2. Lionel A. Stange, Division of Plant Industry, Florida Department of Agriculture and Consumer Services, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.

1'. Body predominately black with first three abdominal segments with large yellow spots; legs, mesoscutellum, and part of vertex rust-orange color *Sphecius speciosus* (Drury)

***Sphecius hogardii* (Latreille)**

This is a Caribbean species known from the Florida Keys (Key West). It has also been recorded in Lover's Key, Lee County, Florida, on 16 July 1999, by collectors C. Porter and L. Stange. It appears to be uncommon in Florida, and nothing is known of its host preferences. The rust-orange coloration is distinctive (Figure 1). However, the black on the abdomen is variable. The Bahama populations lack the black coloration (subspecies *bahamas* Krombein).



Figure 1. *Sphecius hogardii* (Latreille), a cicada killer wasp.
Credits: Division of Plant Industry

***Sphecius speciosus* (Drury)**

This wasp is widely distributed east of the Rocky Mountains and south into Mexico. Sometimes it is found locally in Florida, especially during the summer months when many wasps may nest together gregariously. Human encounters with this giant wasp are only seemingly dangerous since usually it is the stingless male that is the aggressor. The four-foot burrows may have more than 16 cells, which contain one to two cicadas each of the genus *Tibicen*. The wasps are parasitized by sarcophagid flies *Senotainia trilineata* (Wulp); *Metopia argyrocephala* (Meigen); and possibly mutillid wasps.



Figure 2. *Sphecius speciosus* (Drury), a cicada killer wasp.
Credits: Division of Plant Industry

Management

Cicada killers are usually considered beneficial insects since they destroy plant-feeding cicadas. Also, they rarely sting except when the females are handled. However, under certain circumstances, such as when elderly persons or young children are present in the breeding areas, one may want to discourage their presence. This can be done by eliminating or reducing the breeding area, which usually consists of exposed, sandy soil. This area can be mulched or covered with grass. Labeled insecticides can be applied to the nesting sites to kill the wasps.

Selected References

- Bohart RM, Stange LA. 1976. "*Liogorytes joergenseni* (Brèthes), a cicada killer in Argentina." *Pan-Pacific Entomologist* 52: 313.
- Bohart RM, Menke AS. 1976. *Sphecid wasps of the world—a generic revision*. University of California Press. 695 p.
- Dambach CA, Good E. 1943. "Life history and habits of the cicada killer (*Sphecius speciosus*) in Ohio." *Ohio Journal of Science* 43: 32–41.
- Davis WT. 1920. "Mating habits of *Sphecius speciosus*, the cicada killing wasp." *Bulletin Brooklyn Entomological Society* 15: 128–129.
- Eiseley L. 1976. "The coming of the giant wasps." *Defenders of Wildlife*. pp. 152–157.
- Evans HE. 1966. *The comparative ethology and evolution of the sand wasps*. Harvard University Press, Cambridge, xvi + 526 p.

Evans HE, O'Neill KM. 2007. *The sand wasps: Natural history and behavior*. Harvard, Cambridge, Mass. 340 pp.

Holliday CW, Coelho JR. 2006. "Improved key to New World species of (Hymenoptera: Crabronidae)." *Annals of the Entomology Society of America* 99: 793–798.

Holliday CW. (2009). [Prof. Chuck Holliday's Cicada Killer Page](#). Lafayette College. (9 August 2012).

Lin N. 1963. "Territorial behaviour in the cicada killer wasp, *Sphecius speciosus* (Drury). I." *Behaviour* 20: 115–133.

Lin N, Michener CD. 1972. "Evolution of sociality in insects." *Quarterly Review of Biology* 47: 131–159.