

# Parthenium

*Parthenium hysterophorus*



Parthenium is a vigorous species that colonises weak pastures with sparse ground cover. It will readily colonise disturbed, bare areas along roadsides and heavily stocked areas around yards and watering points. Parthenium can also colonise brigalow, gidgee and softwood scrub soils. Its presence reduces the reliability of improved pasture establishment and reduces pasture production potential.

Parthenium is also a health problem as contact with the plant or the pollen can cause serious allergic reactions such as dermatitis and hay fever.

Parthenium is recognised as a Weed of National Significance.

## Legal requirements

Parthenium is a category 3 restricted invasive plant under the *Biosecurity Act 2014*. It must not be given away, sold, or released into the environment. The Act requires everyone to take all reasonable and practical steps to minimise the risks associated with invasive plants under their control. This is called a general biosecurity obligation (GBO). This fact sheet gives examples of how you can meet your GBO.



At a local level, each local government must have a biosecurity plan that covers invasive plants in its area. This plan may include actions to be taken on certain species. Some of these actions may be required under local laws. Contact your local government for more information.

## Description

Parthenium is an annual herb with a deep tap root and an erect stem that becomes woody with age. As it matures, the plant develops many branches in its top half and may eventually reach a height of 2 m.

Its leaves are pale green, deeply lobed and covered with fine soft hairs.

Small creamy white flowers occur on the tips of the numerous stems. Each flower contains four to five black seeds that are wedge-shaped, two millimetres long with two thin, white scales.

## Life cycle

Parthenium normally germinates in spring and early summer, produces flowers and seed throughout its life and dies around late autumn. However, with suitable conditions (rain, available moisture, mild temperatures), parthenium can grow and produce flowers at any time of the year. In summer, plants can flower and set seed within four weeks of germination, particularly if stressed.

## Methods of spread

Parthenium seeds can spread via water, vehicles, machinery, stock, feral and native animals and in feed and seed. Drought conditions aid the spread of seed with increased movements of stock fodder and transports.

## Habitat and distribution

Parthenium is capable of growing in most soil types but becomes most dominant in alkaline, clay loam soils.

The plant is well established in Central Queensland and present in isolated infestations west to Longreach and in northern and southern Queensland.

Infestations have also been found in northern and central parts of New South Wales and it is capable of growing in most states of Australia.

## Control

### Managing parthenium

The GBO requires a person to take reasonable and practical steps to minimise the risks posed by parthenium. This fact sheet provides information and some options for controlling parthenium.

### Prevention and weed seed spread

Pastures maintained in good condition, with high levels of grass crown cover, will limit parthenium colonisation. Drought, and the subsequent reduced pasture cover, creates the ideal window of opportunity for parthenium colonisation when good conditions return.

Vehicles and implements passing through parthenium infested areas should be washed down with water.

Particular care should be taken with earthmoving machinery and harvesting equipment. The wash down procedure should be confined to one area, so that plants that establish from dislodged seed can be destroyed before they set seed.

Extreme caution should be taken when moving cattle from infested to clean areas. Avoid movement during wet periods as cattle readily transport seed in muddy soil. On arrival, cattle should be held in yards or small paddocks until seed has dropped from their coats and tails prior to their release into large paddocks. Infestations around yards can be easily spotted and controlled whereas infestations can develop unnoticed in large paddocks.

Particular care should be taken when purchasing seed, hay and other fodder materials. Always keep a close watch for the emergence of parthenium or other weeds on areas where hay has been fed out.

Property hygiene is important. Owners of clean properties should ensure that visitors from infested areas do not drive through their properties. If your property has parthenium on it, ensure that it is not spread beyond the boundary or further within the property.

### Manual control

Hand pulling of small areas is not recommended. There is a health hazard from allergic reactions and a danger that mature seeds will drop off and increase the area of infestation.

### Pasture management

Grazing management is the most useful method of controlling large-scale parthenium infestations. Maintain pastures in good condition with high levels of ground and grass crown cover. This may require rehabilitation of poor pastures, followed by a sound grazing maintenance program.

**Sown pasture establishment**—Poor establishment of sown pastures can allow parthenium colonisation.

**Pasture agronomy**—Aerial seeding prior to scrub pulling is normally beneficial.

**Overgrazing**—High grazing pressure caused by drought or high stock numbers decreases the vigour and competitiveness of pastures and allows the entry and spread of parthenium. Maintenance of correct stock numbers is most important in controlling parthenium.

**Pastures spelling**—In situations of serious infestation, pasture spelling is essential for rehabilitation. Total spelling is much more effective than simply reducing the stocking rate. However, overgrazing of the remainder of the property must be avoided.

The most appropriate time for pasture spelling is the spring–summer growing period, with the first 6–8 weeks being particularly important. If the condition of perennial grasses (native or sown) is low, spelling for the entire growing season may be required or introduced grasses may need to be re-sown. Herbicide treatment can hasten the rehabilitation process by removing a generation of parthenium seedlings and allowing grass seedlings to establish without competition. In the presence of parthenium, grass establishment is poor.

Grazing during winter should not increase the parthenium risk. Most tropical grasses are dormant and can tolerate moderate grazing during this period. However, parthenium may germinate and grow at this time.

**Fencing**—One of the main problems in controlling parthenium is the large paddock size and the variability of country within paddocks. The resulting uneven grazing pressures encourage parthenium to colonise the heavily grazed country. Ideally, similar land types should be fenced as single units. Fencing can be used to great effect to break up large paddocks, allowing more flexible management such as pasture spelling or herbicide application, options not available previously.

**Burning**—Burning is not promoted as a control strategy for parthenium. However, research suggests that burning for pasture management (e.g. woody weed control) should not result in an increased infestation if the pasture is allowed to recover prior to the resumption of grazing. Stocking of recently burnt areas known or suspected to contain parthenium decreases pasture competition and favours parthenium, ultimately creating a more serious infestation.

## Biological control

The combined effects of biological control agents reduced the density and vigour of parthenium and increased grass production.

There are currently a number of insect species and two rust pathogens that have been introduced to control parthenium—a selection of these are outlined below. *Epiblema strenuana* is a moth introduced from Mexico established in all parthenium areas. The moth's larvae feed inside the stem, forming galls that stunt the plant's growth, reduce competitiveness and seed production.

*Listronotus setosipennis* is a stem-boring weevil from Argentina but is of limited success in reducing parthenium infestations.

*Zygogramma bicolorata* is a defoliating beetle from Mexico which is highly effective where present. It emerges in late spring and is active until autumn.

*Smicronyx lutulentus* (Mexico) lays eggs in the flower buds where the larvae feed on the seed heads. *Conotrachelus albocinereus* (stem-galling weevil from Argentina) produces small galls and is still becoming established in Queensland.

*Bucculatrix parthenica* (leaf mining moth from Mexico) larvae feed on leaves, leaving clear windows in the leaf. *Carmentia ithacae* is a stem boring moth from Mexico which is becoming established at favourable sites in the northern Central Highlands.

*Puccinia abrupta* is a winter rust from Mexico that infects and damages leaves and stems. It is currently established over a wide area from Clermont south. It requires a night temperature of less than 16° and 5–6 hours of leaf wetness (dew). Sporadic outbreaks occur where weather conditions are suitable.

*Puccinia melampodii* is a summer rust from Mexico that weakens the plant by damaging the leaves over the summer growing season. It is currently established and spreading at a number of sites from north of Charters Towers to Injune in the south.

## Herbicide control

### Non-crop areas

Parthenium should be sprayed early before it can set seed. A close watch should be kept on treated areas for at least two years.

Small and/or isolated infestations should be treated immediately. Herbicide control will involve a knockdown herbicide to kill plants that are present and a residual herbicide to control future germinations. Repeated spraying may be required even within the one growing season to prevent further seed production.

Extensive infestations will require herbicide treatment in conjunction with pasture management. Timing of spraying is critical so that parthenium is removed when plants are small and before seeding has occurred. Grasses should be actively growing and seeding so that they can recolonise the infested area.

Table 1 details the herbicides registered for parthenium control and application rates. All herbicides must be applied strictly in accordance with the directions on the label.

### Cropping areas

Controlling parthenium in cropland requires selective herbicide use and/or crop rotations. For further information on parthenium control in crops consult your local biosecurity officer.

## Further information

Further information is available from your local government office, or by contacting Biosecurity Queensland on 13 25 23 or visit [biosecurity.qld.gov.au](http://biosecurity.qld.gov.au).



**Table 1. Herbicides for the control of parthenium**

Situation	Herbicide	Rate	Comments
Pastures, rights-of-way and industrial land	2,4-D as amine 625 g/L (e.g. Ken-Amine 625)	320 mL/100 L water	Spot spray Apply to young actively growing plants, ensuring thorough coverage
	2,4-D as amine 700 g/L (e.g. Amicide Advance 700)	285 mL/100 L water	
Non agricultural areas (native pastures), commercial and industrial areas and rights-of-way	Aminopyralid 375 g/kg plus Metsulfuron-methyl 300 g/kg (Stinger)	10 g/100 L water plus wetting agent (consult label)	Spray to thoroughly wet all foliage but not to cause run-off
Fields and fallow, various crops (consult label)	Atrazine 500 g/L (e.g. Kenso Atrazine 500)	3.6–6 L/ha Rate varies with situation (consult label)	Boom spray. Pre and post emergent application Restrictions apply (consult label) Max 3 kg a.i./ha/yr
Roadside and rights-of-way		6 L/ha	Boom spray. Pre and post emergent application Restrictions apply (consult label) Max 3 kg a.i./ha/yr
Fields and fallow and various crops (see label)	Atrazine 900 g/kg (e.g. Atradex WG)	2–3.3 kg/ha Rate varies with situation (consult label)	Boom spray. Pre and post emergent application Restrictions apply (consult label) Max 3 kg a.i./ha/yr
Roadside and rights-of-way		3.3 kg/ha	Boom spray. Pre and post emergent application. Restrictions apply (consult label). Max 3 kg a.i./ha/yr
Non-crop areas, commercial and industrial areas, pastures and rights-of-way	2,4-D 300 g/L + Picloram 75 g/L (e.g. Tordon 75-D)	125 mL/100 L	Spot spray during rosette stage Use at least 3000 L/ha in dense infestations (consult label)
		3 L/ha	Boom spray during rosette stage (consult label)
Native pastures, rights-of-way, commercial and industrial land	Metsulfuron-methyl 600g/L (e.g. Associate)	5 g/100 L water + wetter	Hand gun. Spray to thoroughly wet all foliage but not to cause runoff
		7 g/ha + wetter	Boom spray. For pastures only Treat in rosette stage (consult label)
		5–7 g/h	Boom spray. Lower rate up to 4-leaf stage, higher rate 4-leaf stage to rosette
Wheat, barley, triticale and cereal rye			
Native pastures, rights-of-way, commercial and industrial land	Triclopyr 75 g/L + Metsulfuron-methyl 28 g/L (e.g. Zelam Brush Weed)	125 mL/100 L water	Spot spray plants from rosette to flowering (consult label)
Commercial and industrial areas, rights-of-way, around agricultural buildings	Hexazinone 750 g/kg (e.g. Velpar DF)	1 kg/ha 2 g/10 L/20 m <sup>2</sup>	Boom spray or spot spray
Around agricultural buildings		3.5 L/ha or 7 L/10 L/20 m <sup>2</sup>	
Grass pastures, fallows, various crop and non-crop situations (consult label)	Dicamba 500 g/L (e.g. Kamba 500) Dicamba 700 g/kg	Rates vary with situation (consult label)	Boom spray or spot spray (consult label)

A number of the listed herbicides are available as different formulations, but some may not be registered for parthenium. Check the label for registration, rate and critical comments. Only use products that list parthenium on the label. The registered rates are for non-crop uses. Consult label for in-crop recommendations. For power hand spray or knapsack use, spray plants to the point of runoff.

**Read the label carefully before use. Always use the herbicide in accordance with the directions on the label.**



This fact sheet is developed with funding support from the Land Protection Fund.

Fact sheets are available from Department of Agriculture and Fisheries (DAF) service centres and our Customer Service Centre (telephone 13 25 23). Check our website at [biosecurity.qld.gov.au](http://biosecurity.qld.gov.au) to ensure you have the latest version of this fact sheet. The control methods referred to in this fact sheet should be used in accordance with the restrictions (federal and state legislation, and local government laws) directly or indirectly related to each control method. These restrictions may prevent the use of one or more of the methods referred to, depending on individual circumstances. While every care is taken to ensure the accuracy of this information, DAF does not invite reliance upon it, nor accept responsibility for any loss or damage caused by actions based on it.