

AZ1133 Revised 08/14

WATER USE IN VEGETABLES—CAULIFLOWER ARIZONA WATER SERIES No. 27

Edward C. Martin, Donald C. Slack, E. James Pegelow

Introduction

Arizona grows approximately 10% of the nation's cauliflower. Most of the cauliflower grown in Arizona is grown in Yuma County. As with most crops grown in Arizona, irrigation plays an important role in determining crop yield and quality. This is especially true for vegetable crops such as cauliflower where water stress can translate to poor yields (tonnage) or poor quality (firmness, head size, color).

Commercial Irrigation Management

Most cauliflower in the state is irrigated using furrow irrigation. Some growers utilize drip irrigation, but high start-up costs make it difficult for most growers to switch. However, it is quite common for growers to use sprinklers for germination. The use of sprinklers helps to reduce the amount of water needed for germination and reduces the potential for salts to enter the seed row and cause emergence problems. Cauliflower is moderately sensitive to salt water; therefore, an ECe of 1.3 or less should be used for irrigation.

Cauliflower should be kept stress free throughout the growing cycle. The critical period for cauliflower is during head development. An irrigation threshold of 40% soil water deficit should be targeted to avoid water stress. If using tensiometers or resistance blocks, -40 to -60 centibars should be used to trigger irrigation. Set tensiometer or resistance blocks at a depth of 12 in. This will give an accurate representation of the moisture level in the plant's rootzone.

Water Use by Cauliflower

Two graphs are given to help determine water use by cauliflower throughout the growing season. Figure 1 shows the consumptive use of cauliflower as a function of Heat Units After Planting (HUAP). The temperatures used to develop this curve were 40°F for the lower threshold and 76°F for the upper threshold. The heat units should be calculated using the sinusoidal approach developed by Snyder (1985). Information on daily maximum and minimum temperatures can be obtained from AZMET. Total heat units required for cauliflower are about 2600. Grown mostly during the winter




Figure 1. Average daily water use based on Heat Unit After Planting (HUAP) for cauliflower for a September 15th planting date.




Figure 2. Average daily water use for cauliflower utilizing a calendar schedule for a September 15th planting date.

months, this translates to a growing season of approximately three to four months, depending on the location and time of planting. Caulifl er uses approximately 17 in. of water in a season.

When determining irrigation water needs, do not forget to incorporate the irrigation system's efficiency. For example, if the irrigation system has an efficiency of 75% and the crop requirement is 2 in., you should apply 2.7 in. to make up for the system's inefficiency (2 in./0.75).

Figure 2 shows the average daily water use for cauliflower in central Arizona (Martin et al., 2001). Maximum water use by cauliflower is about 0.17 in. of water per day. This can vary depending on location and time of planting. Peak usage occurs toward the end of the growing cycle, just prior to harvest. The rootzone calculation used for water management in cauliflower is 1.5 ft., although cauliflower roots go deeper.

Reference

AZMET, Arizona Meteorological Network on the web at http://ag.arizona.edu/azmet/

Martin, E.C., A.S. de Oliveira, A.D. Folta, E.J. Pegelow and D.C. Slack. 2001. Development and testing of a small weighing lysimeter system to assess water use in shallow rooted crops. Transactions of the ASAE. 44(1):71-78.

Snyder, R.L. 1985. Hand calculating degree days. Agric. Forest Meteorol., 35: 353-358.

For additional information, contact your local Cooperative Extension office.



THE UNIVERSITY OF ARIZONA COLLEGE OF AGRICULTURE AND LIFE SCIENCES TUCSON, ARIZONA 85721

EDWARD C. MARTIN

County Extension Director Maricopa County Cooperative Extension Professor And Irrigation Specialist Department Of Agriculture & Biosystems Engineering

DONALD C. SLACK

Dean's Chair for Excellence, Cecil Miller Families

E. JAMES PEGELOW

Former Senior Research Specialist

CONTACT:

EDWARD C. MARTIN ecmartin@email.arizona.edu

This information has been reviewed by University faculty. extension.arizona.edu/pubs/az1133-2014.pdf

Originally published: 2009

Other titles from Arizona Cooperative Extension can be found at: extension.arizona.edu/pubs

Any products, services or organizations that are mentioned, shown or indirectly implied in this publication do not imply endorsement by The University of Arizona.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Jeffrey C. Silvertooth, Associate Dean & Director, Extension & Economic Development, College of Agriculture Life Sciences, The University of Arizona.

The University of Arizona is an equal opportunity, affirmative action institution. The University does not discriminate on the basis of race, color, religion, sex, national origin, age, disability, veteran status, or sexual orientation in its programs and activities.