



## Pacific Pests and Pathogens - Fact Sheets

### Biocontrols - *Diadegma* (285)



Photo 1. *Diadegma* seeking its host.



Photo 2. *Diadegma* has located a second stage diamondback moth caterpillar.



Photo 3. *Diadegma* laying an egg in a second stage diamondback moth caterpillar.



Photo 4. *Diadegma* wasp preening or cleaning its body after egg laying.



Photo 5. The pupa of diamondback moth, yellowish and about 8 mm long. It is blunt at one end and tapered at the other.



Photo 6. Pupa of *Diadegma*. The pupa if the diamondback moth is destroyed by *Diadegma*, except for the silken cocoon. The *Diadegma* cocoon is about the same size as that of the diamondback moth. They are pink/cream, then dark brown.



Photo 7. Pupa of *Diadegma* becomes dark before emergence of the adult. It is elliptical (egg-shaped) not tapered, and the diamondback moth skin is at one end.

#### Common Name

Parasitic wasp

#### Scientific Name

*Diadegma semiclausum*, an ichneumonid (previously known as *Diadegma euerophaga*, *Diadegma xylostellae*, *Angita cerophaga*).

#### Distribution

Worldwide. *Diadegma semiclausum* is native to Europe, but has been introduced into many countries in Asia, Africa and Oceania. It is established in Australia and New Zealand but, it has not established in other Pacific islands, with the exception of the highlands of Papua New Guinea. The wasp has also been introduced into the highlands of

India, Indonesia, Laos, Malaysia, the Philippines, Taiwan, Thailand, and Vietnam. In tropical countries, it does not survive below about 800 m.

## Prey

Diamondback moth, *Plutella xylostella*.

## Symptoms & Life Cycle

*Diadegma* is most commonly found parasitising diamondback moth, although it has been reared from other species. It is a larval parasitoid, preferring to lay its eggs in small larvae, those that have just left their egg masses on *Brassica* leaves.

When the wasp approaches (Photos 1&2), the moth larvae may react with jerky movements, or drop from the leaf on silken threads to escape attack. Those that remain become the target of the wasp which quickly inserts a single egg into each larva (Photos 3&4). The larvae are parasitised once only as the wasp can tell if the diamondback moth larvae have already been attacked.

Parasitised moth larvae develop normally, and outwardly look the same as those that are unparasitised. However, when the moth develops a cocoon (Photo 5), the larva of the wasp makes its own cocoon inside and then develops into a pupa (Photo 6). After 8-10 days the adult emerges.

The life cycle of *Diadegma* is 18-20 days. The female wasp can live for about 25 days if a suitable sugar food is available. During that time it can lay many hundreds of eggs.

The parasitoid prefers temperatures in the range 15-25°C. Therefore, in tropical countries it is most abundant in the cooler highlands.

## Impact

*Diadegma* is a very effective natural enemy of diamondback moth, and very high rates of parasitism (>90%) often occur, provided that broad-spectrum insecticides are not used.

## Detection & Inspection

Look for the small black wasps flying between brassica plants, often hovering near leaves damaged by diamondback moth larvae. Chemicals from the damage attracts them. Look for the wasps searching for diamondback moth larvae on the leaves. Look for *Diadegma* pupae which are formed inside the silken nets spun by the diamondback moths when they pupate. *Diadegma* pupae are oval-shaped, whereas those of diamondback moth are tapered at one end.

## Management as Biocontrol Agents

*Diadegma* is very sensitive to broad-spectrum insecticides, such as pyrethroids and organophosphates. They should not be used if the wasp is active in the crop, otherwise they will kill the wasp and make control of diamondback moth difficult. Instead, if an insecticide is required, use Bt, *Bacillus thuringiensis*, or another selective insecticide. Together, *Diadegma* and Bt can form the basis of an effective IPM program for diamondback moth.

---

AUTHORS Mike Furlong and Grahame Jackson

Photos 1-6 Mike Furlong, University of Queensland, Brisbane, Queensland.

Produced with support from the Australian Centre for International Agricultural Research under project PC/2010/090: *Strengthening integrated crop management research in the Pacific Islands in support of sustainable intensification of high-value crop production*, implemented by the University of Queensland and the Secretariat of the Pacific Community.

---

This fact sheet is a part of the app *Pacific Pests and Pathogens*

The mobile application is available from the Google Play Store and Apple iTunes.



Android Edition



Apple iOS Edition



Australian Government  
Australian Centre for  
International Agricultural Research

---

*Copyright © 2017. All rights reserved.*